



# Childhood vaccinations and when to have them

**8 Weeks**



- ★ **6-in-1** (Diphtheria, Tetanus, Whooping cough, Polio, Hib, Hepatitis B)
- ★ **MenB**
- ★ **Rotavirus** (drops to swallow)

**12 Weeks**



- ★ **6-in-1** (Diphtheria, Tetanus, Whooping cough, Polio, Hib, Hepatitis B)
- ★ **MenB**
- ★ **Rotavirus** (drops to swallow)

**16 Weeks**



- ★ **6-in-1** (Diphtheria, Tetanus, Whooping cough, Polio, Hib, Hepatitis B)
- ★ **Pneumococcal**

**1 Year**



- ★ **MMR** (Measles, Mumps, Rubella)
- ★ **Pneumococcal**
- ★ **MenB**

**3 Years  
4 Months**



- ★ **4-in-1** (Diphtheria, Tetanus, Whooping cough, Polio)
- ★ **MMR** (Measles, Mumps, Rubella)

Getting vaccines on time gives the best protection. If your child misses a dose, contact your GP to catch up. To be fully protected, your child needs all the doses shown.

**Questions about vaccinations?**

**Ask your GP, nurse, health visitor or pharmacist.**

Scan QR  
to find  
out more

